

	Policy:	Dimenhydrinate (Gravol) for Nausea and/or Vomiting
	Number:	
Approved by: MAC November 2009	Manual:	
Signature:	Section:	Medical Directive Emergency
	Page:	1 of 3

Procedure:

Dimenhydrinate (Gravol) 50 mg IM times one dose; or
Dimenhydrinate (Gravol) 25 mg IV times one dose

Indications:

Adult > 16 and

Nausea and/ or Vomiting causing physical distress (discomfort, dehydration) as assessed by a

Registered Nurse; and

Emergency Department out-patient

Contraindications;

Known allergy to Dimenhydrinate; or

Received Dimenhydrinate in the previous 3 hours; or

Altered level of consciousness

Effective Date: November 2009	Revised Date:	Version: November, 2009
File Name: Dimenhydrinate for Adults ED-Med-07		

References:

1. Gay LN, Carlinger PE. The prevention and treatment of motion sickness. I. Seasickness. *Science* 1949; 109: 359. [\[Free Full Text\]](#)
2. Tang J, Watcha MF, White PF. A comparison of costs and efficacy of ondansetron and droperidol as prophylactic antiemetic therapy for elective outpatient gynecologic procedures. *Anesth Analg* 1996; 83: 304–13. [\[Abstract\]](#)
3. Pandit SK, Kothary SP, Pandit UA, et al. Dose-response study of droperidol and metoclopramide as antiemetics for outpatients. *Anesth Analg* 1989; 68: 798–802. [\[Free Full Text\]](#)
4. Grond S, Lynch J, Diefenbach C, et al. Comparison of ondansetron and droperidol in the prevention of nausea and vomiting after inpatient minor gynecologic surgery. *Anesth Analg* 1995; 81: 603–7. [\[Abstract\]](#)
5. Henzi I, Sonderegger J, Tramer M. Efficacy, dose-response and adverse effects of droperidol for prevention of postoperative nausea and vomiting. *Can J Anaesth* 2000; 47: 537–51. [\[Web of Science\]](#)[\[Medline\]](#)
6. Woollorton E. Droperidol: cardiovascular toxicity and deaths [editorial]. *CMAJ* 2002; 166: 932. [\[Free Full Text\]](#)
7. Brown NJ, Roberts LJ. Histamine, Bradykinin and their antagonists. In: Hardman JG, Limbard LL, Gilman AG, eds. *Goodman's and Gilman's the pharmacological basis of therapeutics*. 10th ed. New York: McGraw-Hill, 2001: 645–67.
8. Peterson MC. Letter to the editor. *Anesthesiology* 1949; 10: 767.
9. Wolfe WB. The use of "dramamine" in the prevention of postoperative nausea and vomiting. *Ann Surg* 1952; 136: 261–6. [\[Web of Science\]](#)[\[Medline\]](#)
10. Moore DC, Bridenbaugh LD, Green JC, et al. Intramuscular use of dimenhydrinate (Dramamine) to control postoperative vomiting. *JAMA* 1955; 159: 1342–5.
11. Gordon RA, Vandewater SL, Sleath GE, Caplin D. A study of the value of dimenhydrinate and promethazine hydrochloride in the control of post-anaesthetic vomiting. *Can Anaesth Soc J* 1954; 1: 95–103.

12. Eberhart LHJ, Seeling W, Hartschuh T, et al. Droperidol and dimenhydrinate alone or in combination for the prevention of postoperative nausea and vomiting after nasal surgery in male patients. *Eur J Anaesthesiol* 1999; 16: 790–5. [\[Web of Science\]](#)[\[Medline\]](#)
13. Kothari SN, Boyd WC, Bottcher ML, et al. Antiemetic efficacy of prophylactic dimenhydrinate (Dramamine) versus ondansetron (Zofran): a randomized, prospective trial in patients undergoing laparoscopic cholecystectomy. *Surg Endosc* 2000; 14: 926–9. [\[Web of Science\]](#)[\[Medline\]](#)
14. Kranke P, Morin AM, Roewer N, et al. Dimenhydrinate for prophylaxis of postoperative nausea and vomiting: a meta-analysis of randomized controlled trials. *Acta Anaesthesiol Scand* 2002; 46: 238–44. [\[Web of Science\]](#)[\[Medline\]](#)
15. Tang J, Xiaoguang C, White PF, et al. Antiemetic prophylaxis for office-based surgery: are the 5-HT₃ receptor antagonists beneficial? *Anesthesiology* 2000; 98: 293–8.
16. Gan TJ. Postoperative nausea and vomiting: can it be eliminated? *JAMA* 2002; 287: 1233–6. [\[Free Full Text\]](#)
17. Tramer MR. A rational approach to the control of postoperative nausea and vomiting: evidence from systematic reviews. I. Efficacy and harm of antiemetic interventions, and methodological issues. *Acta Anaesthesiol Scand* 2001; 45: 4–13. [\[Web of Science\]](#)[\[Medline\]](#)
18. Heffernan AM, Rowbotham DJ. Postoperative nausea and vomiting: time for balanced antiemesis? [editorial] *Br J Anaesth* 2000; 85: 675–6. [\[Free Full Text\]](#)
19. Scuderi PE, James RL, Harris L, Mims GR. Multimodal antiemetic management prevents early postoperative vomiting after outpatient laparoscopy. *Anesth Analg* 2000; 91: 1408–14. [\[Abstract/Free Full Text\]](#)