

## EMERGENCY DEPARTMENT PHYSICIAN'S ORDERS

### SUSPECTED RENAL COLIC

Allergies: \_\_\_\_\_

#### 1. Nursing

- Clear Fluids only
- IV NS lock preferably # 18 g
- IV \_\_\_\_\_ @ \_\_\_\_\_ ml/hr
- Vitals q 1 h
- Strain all urine
- Stone for pathology if passed

#### 2. Investigations

- CBC, lytes, urea, creatinine, glucose
- INR, PTT
- Type and screen
- Urine R&M, C&S
- KUB
- CT Pyelogram (Non contrast Abdomen and Pelvis); **caution in patients with known stone disease who lack hydronephrosis on ultrasound and whose pain has been controlled (risk of repeat radiation exposure over time)**
- Move Bedside US machine to bedside (to confirm Aorta Maximum AP diameter < 3cm)
- Ultrasound Renal Imaging Study (KUB)
- \_\_\_\_\_
- \_\_\_\_\_

#### 3. Medications

- Acetaminophen per ED Nursing Directive
- Ibuprofen per ED Nursing Directive
- Indocid Suppository 100 mg PR times one dose
- Ketorolac 30 mg IV times one dose
- Fentanyl per ED nursing directive
- Morphine \_\_\_\_\_ mg IV (suggest 0.1mg/kg)q \_\_\_\_\_ min prn max \_\_\_\_\_ mg
- Gravol \_\_\_\_\_ mg IV (suggest 25-50 mg) q \_\_\_\_\_ hours prn
- Flomax SR 0.4 mg po x one dose
- \_\_\_\_\_
- \_\_\_\_\_

Date \_\_\_\_\_

Physician's Signature \_\_\_\_\_

Time \_\_\_\_\_

Print Name \_\_\_\_\_