

Table 1: Demographics

Variable	App n (%) 52 (49%)	Control n (%) 55 (51%)	P-value
Sex			<i>P=0.68</i>
Male	26 (50%)	25 (45%)	
Female	26 (50%)	30 (55%)	
Age- Median (range)	42 (19-73)	50 (21-86)	<i>P=0.0099</i>
DX			<i>P=0.99</i>
Colon Malignancy	10 (19%)	15 (27%)	
IBD	31 (60%)	34 (62%)	
Other	11 (21%)	6 (12%)	
Immunosuppression			<i>P=0.99</i>
Yes	12 (23%)	12 (22%)	
No	39 (75%)	42 (76%)	
BMI- Median (range)	25 (17-36)	25 (17-46)	<i>P=0.36</i>
Surgical Approach			<i>P=.14</i>
Laparoscopic	31 (60%)	24 (44%)	
Open	19 (36%)	28 (50%)	
Converted	1 (2%)	2 (4%)	
Procedure			<i>P=0.22</i>
Ileocolic resection	7 (13%)	9 (16%)	
Right hemicolectomy	6 (11%)	2 (4%)	
Left hemi	0 (0%)	2 (4%)	
Sigmoid colectomy	2 (4%)	2 (4%)	
Total colectomy	9 (17%)	10 (18%)	
APR	2 (4%)	3 (5%)	
LAR	4 (8%)	5 (9%)	
IPAA	5 (10%)	2 (4%)	
Ileostomy reversal	3 (6%)	6 (11%)	
Other	14 (27%)	14 (25%)	
Stoma			<i>P=0.85</i>
Ileostomy	19 (37%)	19 (35%)	
Colostomy	4 (7%)	5 (9%)	
No stoma	29 (56%)	31 (56%)	